

## References

- [1] D.W. Bartlett, The role of erosion in tooth wear: aetiology, prevention and management, *Int. Dent. J.* 55 (2005) 277–284.
- [2] M.K. Al-Omiri, P.J. Lamey, T. Clifford, Impact of tooth wear on daily living, *Int. J. Prosthodont.* 19 (2006) 601–605.
- [3] C. Ganss, Definition of erosion and links to tooth wear, *Monogr. Oral Sci.* 20 (2006) 9–16.
- [4] D.L. Gambon, H.S. Brand, E.C. Veerman, Dental erosion in the 21st century: what is happening to nutritional habits and lifestyle in our society? *Br. Dent. J.* 213 (2012) 55–57.

- [5] S. Taji, W.K. Seow, A literature review of dental erosion in children, *Aust. Dent. J.* 55 (2010) 358–367.
- [6] A. Johansson, A.K. Johansson, R. Omar, G.E. Carlsson, Rehabilitation of the worn dentition, *J. Oral Rehabil.* 35 (2008) 548–566.
- [7] M. Fradeani, G. Barducci, L. Bacherini, M. Brennan, Esthetic rehabilitation of a severely worn dentition with minimally invasive prosthetic procedures (MIPP), *Int. J. Periodont. Restor. Dent.* 32 (2012) 135–147.
- [8] I.A. Meyers, Minimum intervention dentistry and the management of tooth wear in general practice, *Aust. Dent. J.* 58 (Suppl. 1) (2013) 60–65.
- [9] L. Sabahipour, D. Bartlett, A questionnaire based study to investigate the variations in the management of tooth wear by UK and prosthodontists from other countries, *Eur. J. Prosthodont. Restor. Dent.* 17 (2009) 61–66.
- [10] B.A. Loomans, M.V. Cardoso, F.J. Roeters, N.J. Opdam, J. De Munck, M.C. Huysmans, et al., Is there one optimal repair technique for all composites? *Dent. Mater.* 27 (2011) 701–709.
- [11] N.J. Opdam, E.M. Bronkhorst, B.A. Loomans, M.C. Huysmans, Longevity of repaired restorations: a practice based study, *J. Dent.* 40 (2012) 829–835.
- [12] P.A. Da Rosa Rodolpho, T.A. Donassollo, M.S. Cenci, A.D. Loguercio, R.R. Moraes, E.M. Bronkhorst, et al., 22-Year clinical evaluation of the performance of two posterior composites with different filler characteristics, *Dent. Mater.* 27 (2011) 955–963.
- [13] K.E. Ahmed, S. Murbay, Survival rates of anterior composites in managing tooth wear: systematic review, *J. Oral Rehabil.* (2015).
- [14] J.P.T. Higgins, S. Green, *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 [updated March 2011], The Cochrane Collaboration, London, 2011.
- [15] A. Liberati, D.G. Altman, J. Tetzlaff, C. Mulrow, P.C. Gotzsche, J.P. Ioannidis, et al., The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration, *J. Clin. Epidemiol.* 62 (2009) e1–34.
- [16] K.A. Robinson, K. Dickersin, Development of a highly sensitive search strategy for the retrieval of reports of controlled trials using PubMed, *Int. J. Epidemiol.* 31 (2002) 150–153.
- [17] A.S. Al-Khayatt, A. Ray-Chaudhuri, N.J. Poyser, P.F. Briggs, R.W. Porter, M.G. Kelleher, et al., Direct composite restorations for the worn mandibular anterior dentition: a 7-year follow-up of a prospective randomised controlled split-mouth clinical trial, *J. Oral Rehabil.* 40 (2013) 389–401.
- [18] A.B. Gulamali, K.W. Hemmings, C.J. Tredwin, A. Petrie, Survival analysis of composite Dahl restorations provided to manage localised anterior tooth wear (ten year follow-up), *Br. Dent. J.* 211 (2011) E9.
- [19] D. Bartlett, G. Sundaram, An up to 3-year randomized clinical study comparing indirect and direct resin composites used to restore worn posterior teeth, *Int. J. Prosthodont.* 19 (2006) 613–617.
- [20] A.M. Gow, K.W. Hemmings, The treatment of localised anterior tooth wear with indirect Artglass restorations at an increased occlusal vertical dimension: results after two years, *Eur. J. Prosthodont. Restor. Dent.* 10 (2002) 101–105.
- [21] K.W. Hemmings, U.R. Darbar, S. Vaughan, Tooth wear treated with direct composite restorations at an increased vertical dimension: results at 30 months, *J. Prosthet. Dent.* 83 (2000) 287–293.
- [22] P.R. Schmidlin, T. Filli, C. Imfeld, S. Tepper, T. Attin, Three-year evaluation of posterior vertical bite reconstruction using direct resin composite—a case series, *Oper. Dent.* 34 (2009) 102–108.
- [23] A.W. Walls, The use of adhesively retained all-porcelain veneers during the management of fractured and worn anterior teeth: part 2. Clinical results after 5 years of follow-up, *Br. Dent. J.* 178 (1995) 337–340.
- [24] N.J. Poyser, R.W. Porter, P.F. Briggs, H.S. Chana, M.G. Kelleher, The Dahl Concept: past, present and future, *Br. Dent. J.* 198 (2005) 669–676 (quiz 720).
- [25] T. Attin, T. Filli, C. Imfeld, P.R. Schmidlin, Composite vertical bite reconstructions in eroded dentitions after 5.5 years: a case series, *J. Oral Rehabil.* 39 (2012) 73–79.
- [26] F.S. Nohl, P.A. King, K.E. Harley, Ibbetson RJ: Retrospective survey of resin-retained cast-metal palatal veneers for the treatment of anterior palatal tooth wear, *Quintessence Int.* 28 (1997) 7–14.
- [27] R.J. Smales, T.L. Berekally, Long-term survival of direct and indirect restorations placed for the treatment of advanced tooth wear, *Eur. J. Prosthodont. Restor. Dent.* 15 (2007) 2–6.
- [28] J.T. Hamburger, N.J. Opdam, E.M. Bronkhorst, C.M. Kreulen, J.J. Roeters, M.C. Huysmans, Clinical performance of direct composite restorations for treatment of severe tooth wear, *J. Adhes. Dent.* 13 (2011) 585–593.
- [29] F. Vailati, L. Gruetter, U.C. Belsler, Adhesively restored anterior maxillary dentitions affected by severe erosion: up to 6-year results of a prospective clinical study, *Eur. J. Esthet. Dent.* 8 (2013) 506–530.
- [30] N.J. Poyser, P.F. Briggs, H.S. Chana, M.G. Kelleher, R.W. Porter, M.M. Patel, The evaluation of direct composite restorations for the worn mandibular anterior dentition—clinical performance and patient satisfaction, *J. Oral Rehabil.* 34 (2007) 361–376.
- [31] R. Hickel, A. Peschke, M. Tyas, I. Mjor, S. Bayne, M. Peters, et al., FDI World Dental Federation—clinical criteria for the evaluation of direct and indirect restorations: update and clinical examples, *J. Adhes. Dent.* 12 (2010) 259–272.
- [32] J.F. Cvar, G. Ryge, Reprint of criteria for the clinical evaluation of dental restorative materials. 1971, *Clin. Oral Investig.* 9 (2005) 215–232.
- [33] N.J. Opdam, E.M. Bronkhorst, M.S. Cenci, M.C. Huysmans, N.H. Wilson, Age of failed restorations: a deceptive longevity parameter, *J. Dent.* 39 (2011) 225–230.
- [34] T. Laegreid, N.R. Gjerdet, A.K. Johansson, Extensive composite molar restorations: 3 years clinical evaluation, *Acta Odontol. Scand.* 70 (2012) 344–352.
- [35] N.J. Opdam, F.H. van de Sande, E. Bronkhorst, M.S. Cenci, P. Bottenberg, U. Pallesen, et al., Longevity of posterior composite restorations: a systematic review and meta-analysis, *J. Dent. Res.* 93 (2014) 943–949.
- [36] F.H. van de Sande, P.A. Rodolpho, G.R. Basso, R. Patias, Q.F. da Rosa, F.F. Demarco, et al., 18-year survival of posterior composite resin restorations with and without glass ionomer cement as base, *Dent. Mater.* 31 (2015) 669–675.
- [37] M.B. Correa, M.A. Peres, K.G. Peres, B.L. Horta, A.J. Barros, F.F. Demarco, Do socioeconomic determinants affect the quality of posterior dental restorations? A multilevel approach, *J. Dent.* 41 (2013) 960–967.
- [38] F.H. van de Sande, N.J. Opdam, P.A. Rodolpho, M.B. Correa, F.F. Demarco, M.S. Cenci, Patient risk factors' influence on survival of posterior composites, *J. Dent. Res.* 92 (2013) 78S–83S.
- [39] J.W. van Dijken, U. Pallesen, Fracture frequency and longevity of fractured resin composite, polyacid-modified resin composite, and resin-modified glass ionomer cement class IV restorations: an up to 14 years of follow-up, *Clin. Oral Investig.* 14 (2010) 217–222.
- [40] D. Hurst, What is the best way to restore the worn dentition? *Evid. Based Dent.* 12 (2011) 55–56.
- [41] M. Dorri, All-ceramic tooth-supported single crowns have acceptable 5-year survival rates, *Evid. Based Dent.* 14 (2013) 47.
- [42] M. Kimmich, C.F. Stappert, Intraoral treatment of veneering porcelain chipping of fixed dental restorations: a review and clinical application, *J. Am. Dent. Assoc.* 144 (2013) 31–44.
- [43] T.E. Donovan, Commentary: evaluation of fracture resistance and failure risks of posterior partial coverage restorations, *J. Esthet. Restor. Dent.* 25 (2013) 123–124.
- [44] J.F. Guth, E.S.J.S. Almeida, M. Ramberger, F. Beuer, D. Edelhofer, Treatment concept with CAD/CAM-fabricated high-density polymer temporary restorations, *J. Esthet. Restor. Dent.* 24 (2012) 310–318.